

Line bundles in supersymmetric coset models

S. Groot Nibbelink

NIKHEF, P.O. Box 41882

1009 DB, Amsterdam, The Netherlands

Revised version

Abstract

The scalars of an $N = 1$ supersymmetric σ -model in 4 dimensions parameterize a Kähler manifold. The transformations of their fermionic superpartners under the isometries are often anomalous. These anomalies can be canceled by introducing additional chiral multiplets with appropriate charges. To obtain the right charges a non-trivial singlet compensating multiplet can be used. However when the topology of the underlying Kähler manifold is non-trivial, the consistency of this multiplet requires that its charge is quantized. This singlet can be interpreted as a section of a line bundle. We determine the Kähler potentials corresponding to the minimal non-trivial singlet chiral superfields for any compact Kählerian coset space G/H . The quantization condition may be in conflict with the requirement of anomaly cancelation. To illustrate this, we discuss the consistency of anomaly free models based on the coset spaces $E_6/SO(10) \times U(1)$ and $SU(5)/SU(2) \times U(1) \times SU(3)$.

PACS: 02.20.Sv, 02.40.-k, 12.60.Jv, 11.30.Na.

Keywords: Supersymmetry, coset space, Kähler potential, line bundle, anomaly cancelation.

Non-linear σ -models in 4 dimensions with $N = 1$ supersymmetry may be the effective field theory setting for the discussion of models beyond the standard model. Chiral fermions ψ_L^α are contained in an irreducible representation of $N = 1$ supersymmetry called chiral multiplets $\Phi^\alpha = (z^\alpha, \psi_L^\alpha, h^\alpha)$. The lowest components of those multiplets z^α are complex scalars that are the coordinates of a complex Kähler manifold $\mathcal{M}$ [1]. Locally the metric $g_{\alpha\bar{\alpha}}$ of this manifold is obtained from a Kähler potential $K(z, \bar{z})$. Non-trivial examples of Kähler manifolds are provided by homogeneous coset spaces G/H , where G is any compact Lie group and H is the centralizer of a torus in G .

The chiral fermion content of a supersymmetric model based on the Kähler manifold $\mathcal{M}$ is often anomalous [2, 3]. This leads to the conclusion that supersymmetric models based on G/H cosets are inconsistent unless a complete mirror sector is introduced, which is phenomenologically uninteresting as it easily results in large fermion masses [4, 5]. However one can avoid this by including additional non-mirror (matter) chiral multiplets to cancel anomalies. In order that these matter multiplets respect all isometries of $\mathcal{M}$ they can be introduced as tensor products of co(ntra)variant vectors on $\mathcal{M}$ [9, 10]. This type of matter alone is often not sufficient to build anomaly free models: the charge assignment is too restrictive. To overcome this difficulty, a singlet chiral multiplet was introduced in ref. [6], that transforms non-trivially under the isometries of $\mathcal{M}$. This non-trivial singlet superfield can be used as a compensator to give physical chiral multiplets the charges required for isometry anomaly cancellation. For this non-trivial singlet multiplet, as well as for the other matter multiplets, invariant Kähler potentials have to be constructed. From these one obtains a supersymmetric lagrangean, that is invariant under the action of the isometries of $\mathcal{M}$.

The definitions of matter chiral superfields, as given above, take only the local properties of the Kähler manifold $\mathcal{M}$ into account. In order to guarantee that the definitions of these matter multiplets do not lead to global inconsistencies, additional constraints have to be imposed. In mathematics a well-defined function on a manifold is called section of a bundle over this manifold. [7, 8]. In this language, a globally well-defined matter multiplet has a scalar component that is a section of a bundle. A co(ntra)variant vector on $\mathcal{M}$ can be interpreted globally as a section of the (co)tangent bundle and is therefore well-defined. The global definition of the non-trivial singlet compensating multiplet is more involved: when the topology of the Kähler manifold $\mathcal{M}$ is non-trivial, there is an additional quantization condition [7], called the cocycle condition. The corresponding bundle is called a complex line bundle. This quantization condition is equivalent to the requirement, that the integral over the Kähler form associated with the Kähler potential of the non-trivial singlet is equal to $2\pi\mathbb{Z}$. In particular there is a minimal charge; all charges of non-trivial singlets have to be an integer number times the minimal charge. This condition leads to quantization of Newton's constant in supergravity theories [11],

when the Kähler potential is covariant but not invariant. The condition also restricts or can even be in conflict with the freedom of the charge assignment which was used in refs. [6, 12] to obtain anomaly free supersymmetric models based on the coset G/H .

The objectives of this letter are the following. We want to determine the minimal non-trivial singlet multiplets that can be coupled to compact Kählerian coset spaces G/H . Such a singlet multiplet can be used in supersymmetric model building to obtain the appropriate charge assignment for anomaly cancelation. To this end we have to identify the minimal line bundles, which can be done by showing that the corresponding Kähler form satisfies a minimal integral condition. To obtain general results, we introduce some Lie group machinery to describe G/H coset spaces for any compact simple Lie group G . Using this we review the construction of Kähler potentials for these homogeneous spaces following refs. [13, 14] with a few minor changes in notation. We obtain a generating set of Kähler potentials K^J that satisfy the minimal cocycle condition and which may be used to construct invariant lagrangeans for non-trivial singlets superfields. The last step is to determine the charges of these superfields. It is essential to know these charges as they may be such that anomaly cancelation is not possible. To show that is a non-trivial requirement, we investigate whether the construction of two anomaly free models is compatible with the cocycle condition: $E_6/SO(10) \times U(1)$ [6] and a Grassmannian version of the standard model [12]. In the latter model matter is introduced by using that the transformation rules of covariant vectors factorizes. For completeness, we check the consistency of these matter representations as well.

We start by discussing the quantization condition of the charge of non-trivial singlet representation when the topology of $\mathcal{M}$ is non-trivial: the cocycle condition for a complex line bundle. By using the covariance of a Kähler potential K

$$K(z', \bar{z}') = K(z, \bar{z}) - c(z) - c^\dagger(\bar{z}) \quad (1)$$

under the isometries of $\mathcal{M}$, the transformation rule

$$s' = e^{-c(z)} s \quad (2)$$

of the scalar component s of a non-trivial chiral multiplet is obtained. The invariant Kähler potential

$$K_{\text{line}}(s, \bar{s}; z, \bar{z}) = \bar{s} e^{-K(z, \bar{z})} s. \quad (3)$$

can be used for supersymmetric model building purposes [6].

A well-defined complex line bundle over a manifold satisfies consistency conditions [7] that insure the global existence of this non-trivial singlet representation. In order that

this singlet can be interpreted as a section of a line bundle, the cocycle condition on three overlapping coordinate patches $(1, 2, 3)$ of $\mathcal{M}$ demands that

$$c^{(123)} \equiv c^{(12)} + c^{(23)} + c^{(31)} = -2\pi i \mathbb{Z}. \quad (4)$$

Theorems by de Rham [7, 15] tell us that this requirement is equivalent to the condition that the integral $\frac{1}{2\pi} \int_C \omega(K) = \mathbb{Z}$ of any Kähler 2-form $\omega(K)$ over any 2-cycle C of $\mathcal{M}$ is integer. Locally the Kähler form $\omega(K)$ can be obtained from a Kähler potential K as

$$\omega(K) = -i K_{,\alpha\bar{\alpha}} d\bar{z}^{\alpha} \wedge dz^{\alpha}. \quad (5)$$

In this letter we want to obtain the minimal charges of non-trivial singlet matter couplings to coset spaces G/H . To do this we need to identify a set of Kähler potentials K^J and a set of 2-cycles C_I such that

$$\int_{C_I} \omega(K^J) = 2\pi \delta_I^J. \quad (6)$$

From now on we are only concerned with Kählerian homogeneous coset spaces G/H where G a simple compact Lie group. It is convenient to use the Cartan normalization of the algebra of G which we now review [16, 17, 18]. $T = \{\tau_i\}$ is the set of generators of a Cartan subgroup of G . According to a theorem by Borel [19] a coset G/H is Kählerian if H is the centralizer $C(Y)$ of a torus in G . This torus is generated by the set

$$Y = \{\mathbf{Y}^I = (G^{-1})^{Ii} \tau_i\}. \quad (7)$$

(Hence the index i enumerates the elements of T and I the elements of Y , hence the indices I are a subset of i .) An arbitrary linear combination of τ_i is denoted by τ . Let ϵ_{α} be the generator associated with root α and let $\epsilon_{\pm i} = \epsilon_{\pm \alpha_i}$ denote the creation and annihilation operators of the simple root α_i . Hermitean conjugation of the generators is taken to give $\tau_i^{\dagger} = \tau_i$ and $\epsilon_{\alpha}^{\dagger} = \epsilon_{-\alpha}$. The algebra of G associated with the simple roots can be stated as

$$[\tau_i, \tau_j] = 0, \quad [\epsilon_i, \epsilon_{-j}] = \delta_{ij} \tau_j, \quad [\tau_i, \epsilon_{\pm j}] = \pm G_{ij} \epsilon_{\pm j}, \quad (8)$$

where $G_{ij} \equiv \frac{2\langle \alpha_i, \alpha_j \rangle}{\langle \alpha_i, \alpha_i \rangle} = \alpha_j(\tau_i)$ is the Cartan matrix in this normalization. Following [13, 20, 21] we divided the generators of G into the following sets. We have already defined the set $Y = \{\mathbf{Y}^I\}$. We denote by

$$S = \{S_a\} = \{\tau_i | i \neq I\} \cup \{\epsilon_{\alpha} | \alpha = \sum_{i \neq I} \alpha^i \alpha_i \text{ a root}\} \quad (9)$$

the set of generators of H that are not in Y . The two sets

$$X = \{X_\alpha\} = \{\epsilon_\alpha | \epsilon_\alpha \notin S, \alpha > 0\} \quad \text{and} \quad \bar{X} = \{\bar{X}_\alpha\} = \{\epsilon_\alpha | \epsilon_\alpha \notin S, \alpha < 0\} \quad (10)$$

contain the remaining part of the generators of G .

Using this notation we give representations of elements of H , G/H , etc. An element $h \in H$ is written as $h = e^{i\beta S} e^{i\gamma Y}$, where $\beta^a, \gamma_I \in \mathbb{R}$ and summation over indices has been assumed to be understood. The subset $\hat{H} \subset G^\mathbb{C}$ of the complexification of the group G is generated by Y, S and X . Any element $\hat{h} \in \hat{H}$ of $\hat{H}$ is represented as $\hat{h} = e^{aX} e^{bS} e^{cY}$, where $a^\alpha, b^a, c_I \in \mathbb{C}$ and therefore an element of $G^\mathbb{C}/\hat{H}$ can be written as

$$\xi(z) = e^{z\bar{X}}, \quad (11)$$

where $z^\alpha \in \mathbb{C}$ carries the same root-indices as $\bar{X}_\alpha$. And finally let $U(z, \bar{z})$ denote an element of the unitary representation of G/H . According to [13] $\hat{H}$ is chosen such that $G/H \cong G^\mathbb{C}/\hat{H}$, hence $U(z, \bar{z})$ can be expressed in terms of $\xi(z)$ as

$$U(z, \bar{z}) = \xi(z) e^{A(z, \bar{z})X} e^{B(z, \bar{z})S} e^{-\frac{1}{2}K(z, \bar{z})Y}. \quad (12)$$

The representative $U(z, \bar{z})$ of the equivalence classes of G/H is chosen such that $B(z, \bar{z})$ and $K_J(z, \bar{z})$ are real functions. (The normalization of the functions $K(z, \bar{z})$ is chosen for convenience later on.)

The non-linear transformation properties of the coordinates z and $\bar{z}$ of G/H can be defined using ξ [21, 20] or U [22, 23] by

$$g\xi(z) = \xi(z')\hat{h}(z; g) \quad \text{and} \quad gU(z, \bar{z}) = U(z', \bar{z}')h(z, \bar{z}; g) \quad (13)$$

for any element g of G . The functions $\hat{h}$ and h are chosen such that $\xi(z')$ and $U(z', \bar{z}')$ are again of the forms given above. Combining these transformation rules with the identification of G/H with $G^\mathbb{C}/\hat{H}$ according to eq. (12) shows that

$$K_J(z', \bar{z}') = K_J(z, \bar{z}) - c_J(z; g) - c_J^\dagger(\bar{z}; g); \quad (14)$$

thus the functions $K_J(z, \bar{z})$ transform as Kähler potentials [13]. There it is shown also that the set of Kähler potentials $\{K_J\}$ is complete.

First of all we want to obtain an explicit formula for a fundamental Kähler potential K_J . To this end we consider a representation with orthonormal weight vectors $|\mathbf{b}^J, \mathbf{w}\rangle$ and highest weight $\mathbf{b}^J$, that has all its Dynkin labels zero except for the J th one which is 1: $(b^J)_j = \delta_j^J$. The highest weight vector $|\mathbf{b}^J, \mathbf{b}^J\rangle$ forms a one dimensional irreducible

H -representation, as only generator that acts non-trivially on it (ϵ_{-J}), is not contained in Y or S . Using the BKMU-projector $P^J \equiv |\mathbf{b}^J, \mathbf{b}^J\rangle\langle\mathbf{b}^J, \mathbf{b}^J|$, that projects on this one-dimensional subspace, the fundamental Kähler potentials K^J can be represented as

$$K^J(z, \bar{z}) = K_I(G^{-1})^{IJ} = \ln \det_{P^J} [\xi^\dagger(\bar{z})\xi(z)] . \quad (15)$$

This follows because any Kähler potential $K_{\eta_{\mathbf{b}}}(z, \bar{z})$ obtained using a BKMU-projector $\eta_{\mathbf{b}}$ acting a representation with highest weight $\mathbf{b}$, can be decomposed [13, 14, 21] as

$$K_{\eta_{\mathbf{b}}}(z, \bar{z}) \equiv \ln \det_{\eta_{\mathbf{b}}} [\rho_{\mathbf{b}}(\xi^\dagger(\bar{z}))\rho_{\mathbf{b}}(\xi(z))] = \text{tr}_{\mathbf{b}} \left(\eta_{\mathbf{b}} \tau_I \right) K^I(z, \bar{z}). \quad (16)$$

For the BKMU-projection operators P^J we find that $\text{tr} (P^J \tau_I) = \delta_I^J$, using the properties of the highest weight vector $|\mathbf{b}^J, \mathbf{b}^J\rangle$ and the Cartan matrix.

Next we show that the Kähler forms of the Kähler potentials K^J satisfy the minimal cocycle condition (6). For the generating 2-cycles $C_I : \mathbb{C}P^1 \longrightarrow G^{\mathbb{C}}/\hat{H}$ we take the continuous mappings $v \mapsto (0, \dots, 0, z^{\alpha_I} = v, 0, \dots, 0)$. Using the properties of the projector P^J and the generators $\epsilon_{\pm i}$ we obtain that $K^J|_{C_I}(v, \bar{v}) = \delta_I^J \ln(1 + \bar{v}v)$. This implies that the integral $\int_{C_I} \omega(K^J|_{C_I})$ reduces to the standard integral of the $\mathbb{C}P^1$ -Kähler form over $\mathbb{C}P^1$ itself. As the latter is equal to 2π , we see that the minimal cocycle condition (6) is satisfied.

Finally we determine the charges of the sections of the minimal line bundles, as these are the charges of the non-trivial matter multiplets. The charge of the s^J section of the line bundle is the same as the charge of weight $\mathbf{b}^J$, we find

$$\mathbf{b}^J(\mathbf{Y}^I) = (G^{-1})^{JI}. \quad (17)$$

For the z^{α_J} charge of the coordinates we find

$$\alpha_J(\mathbf{Y}^I) = \delta_J^I. \quad (18)$$

This shows that the minimal $\mathbf{Y}^J$ -charge of a section s^J of a line bundle over G/H is $(G^{-1})^{JI}$ times the charge of z^{α_J} .

We now turn to the consequences of our results for supersymmetric model building. In the construction of a model we may need the transformation rule of a non-trivial compensating singlet to get a $U(1)$ charge assignment such that anomaly cancelation is ensured. However in general it is not guaranteed that the charge we need is an integer multiple of the smallest charge of a non-trivial singlet. How this restriction acts, is now illustrated by two anomaly free models build upon coset spaces.

Ref. [6] discusses an anomaly free model based on the coset $E_6/SO(10) \times U(1)$ [25]. The coset is parameterized by a 16-component spinor of $SO(10)$. To cancel the $U(1)$ -anomaly the model is extended to complete the $\underline{27}$ of E_6 . According to the branching rule

$$\underline{27} = \underline{1}(4) + \underline{16}(1) + \underline{10}(-2), \quad (19)$$

this can be done by introducing a $SO(10)$ singlet with charge 4 and a vector of $SO(10)$ with charge -2 . As discussed in ref. [6] the vector $\underline{10}$ can be obtained from a tensor product of 2 covariant vectors of $E_6/SO(10) \times U(1)$. However this vector $\underline{10}$ of $SO(10)$ has charge 2 and the $SO(10)$ singlet has charge 0. According to the results above the minimal charge of a non-trivial compensating singlet is $G^{JJ} = 4/3$. Hence we need the 3th power of s to define the singlet and a rescaling with the -3 th power of s of the vector to define the vector matter representation with the right charges. Both these powers of s are integers, therefore this models satisfies the line bundle consistency conditions.

Ref. [12] discusses another example of an anomaly free model that is based on the Grassmannian coset $SU(5)/SU(2) \times U(1) \times SU(3)$ [26, 27, 28] with the chiral fermion content of the standard model. For this model the question whether it is globally well-defined is more involved, as matter is not introduced as tangent vectors. To see what the difficulty is, we give a short review of the introduction of matter in this model; for details we refer to ref. [12].

The coordinates Q^{ia} of this coset are interpreted as the superpartners of the left-handed quark doublet q_L^{ia} . The superpartners L and D of a lepton doublet l_L^i and a (down) quark triplet d_L^a were introduced by noticing that the transformation rule for dQ^{ia} factorizes under infinitesimal transformations [12] $\delta dQ = H(Q)dQ + dQ\tilde{H}(Q)$ where $H(Q)$ and $\tilde{H}(Q)$ are holomorphic functions. This can be used to define infinitesimal transformation rules for L and D : $\delta L = H(Q)L$ and $\delta D = -\tilde{H}(Q)D$. Note that this transformation rule is different from the one given in ref. [12], because we $\underline{\bar{3}}$ of $SU(3)$ in stead of the $\underline{3}$ to compile the matter representations of the supersymmetric standard model. As these are only infinitesimal transformations, we do not obtain any information concerning global consistency.

We now show that L and D can be interpreted as sections of bundles over this coset. Using the transformation property (13) of $\xi(Q)$ with a $SU(5)$ matrix given by

$$g = \begin{pmatrix} A & B \\ C & D \end{pmatrix}, \quad (20)$$

where $A \in SU(2)$, $D \in SU(3)$, B a 2×3 and C a 3×2 matrix, we obtain the global transformation rule for Q :

$$Q' = (AQ + B)(D + CQ)^{-1}. \quad (21)$$

By introducing the matrices $\beta = A^{-1}B$ and $\gamma = D^{-1}C$ this can be written as

$$\gamma A^{-1}Q'D = \mathbb{1} - (\mathbb{1} - \gamma\beta)(\mathbb{1} + \gamma Q)^{-1}.$$

From this equation it follows that the transformation of the differential factorizes as

$$\frac{\partial Q'^{ia}}{\partial Q^{jb}} = (G)^i_j (\tilde{G})^a_b, \quad (22)$$

indeed $dQ' = A(\mathbb{1} - \beta\gamma)(\mathbb{1} + Q\gamma)^{-1}dQ(\mathbb{1} + \gamma Q)^{-1}D^{-1}$. Now we know that the tangent bundle is well-defined, therefore the transition functions $G^{(ab)}$ and $\tilde{G}^{(ab)}$ satisfy all consistency conditions by eq. (22). By defining the transition functions acting on L and D via

$$L^{(a)} = G^{(ab)}L^{(b)} \text{ and } D^{(a)} = (\tilde{G}^{(ba)})^{-1}D^{(b)}, \quad (23)$$

well-defined bundles over $SU(5)/SU(2) \times U(1) \times SU(3)$ are obtained.

We now turn to the line bundle constraint. In the normalization of the $SU(5)$ algebra employed in ref. [12], the $U_Y(1)$ charges of Q , L and D are resp. 5, 3 and -2 . According to the results above the charge of a section of the line bundle is $G^{33} = 6/5$ times that of the charge of the coset coordinates Q , therefore the minimum charge is 6. The matching between the weak hypercharge Y_w and the charge Y for the quark doublets requires that $Y = 15Y_w$. It can be checked that with only integer the powers of the minimal line bundle, it is possible to obtain the standard model hyper charges. We conclude that the Grassmannian standard model with non-linear compact $SU(5)$ symmetry can be defined globally.

The main objective of this work was to determine the Kähler potential that satisfy the minimal quantization condition (6) for a general Kählerian compact coset space G/H . The BKMU-projector P^J projects on the highest weight vector of a representation with highest weight $\mathbf{b}^J$ with the Dynkin labels $(b^J)_j = \delta_j^J$. This projector was used to define the Kähler potential K^J (eq. (15)) and its properties showed that this Kähler potential satisfies the minimal cocycle conditions. The minimal charge of a section s^J of a line bundle associated with $\boldsymbol{\tau}_J$ was found to be half of the charge of $z^{\boldsymbol{\alpha}_J}$. In this way we obtained an constructive proof a theorem by Borel-Weil discussed by Serre [24] that classifies the elements of the (co)homology groups of G/H .

The importance of these results for supersymmetric model building lies in the fact that they provide an additional restrictions on the charge assignment of supersymmetric matter coupling and may be in conflict with the requirement of anomaly cancelation. This has been illustrated by the review of two anomaly free models containing a coset space at

their core. The $E_6/SO(10) \times U(1)$ model satisfies the consistency condition. The analysis for $SU(5)/SU(2) \times U(1) \times SU(3)$ was more involved. First we showed that the scalars L^i and D^a are sections of well-defined bundles: using the fact that the transition function of the tangent bundle factorizes, transition functions for these bundles were obtained. We have seen that it is possible to obtain a Grassmannian standard model that satisfies the line bundle constraint as well.

As is clear from these examples the consistent definition of a line bundle provides a stringent restriction on supersymmetric σ -models, in particular those based on coset spaces. With this in mind, it is interesting to study what kind of other supersymmetric models are allowed and what their phenomenology is.

Acknowledgments

I would like to thank C. Hofman for explaining the importance of quantization condition for line bundles to me, J.J. Duistermaat for introducing me to classification of line bundles of G/H coset spaces by the work of Borel and Weil as is discussed in Serre's Bourbaki notes [24] and J.W. van Holten for valuable discussions while this work was in progress.

References

- [1] J.A. Bagger and E. Witten, *Phys. Lett.* **B118** (1982) 103
- [2] G. Moore and P. Nelson, *Phys. Rev. Lett.* **53** (1984) 1519.
- [3] S. Aoyama and J.W. van Holten, *Nucl. Phys.* **B258** (1985) 18.
- [4] W. Buchmuller and W. Lerche, *Ann. Phys.* **175** (1987) 159.
- [5] A. C. W. Kotcheff and G. M. Shore, *Nucl. Phys.* **B333** (1990) 701.
- [6] S. Groot Nibbelink and J. W. van Holten, *Phys. Lett.* **B442** (1998) 185, [hep-th/9808147](#).
- [7] R.O. Wells, Jr, "Differential analysis on complex manifolds," *New York, USA: Springer Verlag* (1980).
- [8] M. Nakahara, "Geometry, Topology And Physics," *Bristol, UK: Hilger* (1990).
- [9] J. W. van Holten, *Nucl. Phys.* **B260** (1985) 125.
- [10] Y. Achiman, S. Aoyama, and J. W. van Holten, *Phys. Lett.* **141B** (1984) 64.

- [11] E. Witten and J. Bagger, *Phys. Lett.* **115B** (1982) 202.
- [12] S. Groot Nibbelink and J.W. van Holten, *hep-th/9903006*, *Nucl. Phys.* **B** (in press).
- [13] K. Itoh, T. Kugo and H. Kunitomo, *Nucl. Phys.* **B263** (1986) 295.
- [14] M. Bando, T. Kugo, and K. Yamawaki, *Phys. Rept.* **164** (1988) 217–314.
- [15] R. Bott and L.W. Tu, “Differential Forms in Algebraic Topology,” *New York, USA: Springer Verlag* (1982).
- [16] R.N. Cahn, “Semisimple Lie Algebras And Their Representations,” *Menlo Park, USA: Benjamin/Cummings* (1984).
- [17] G.G. Bäuerle and E.A. de Kerf, “Lie Algebras. Pt. 1: Finite And Infinite Dimensional Lie Algebras And Applications In Physics,” *Amsterdam, Netherlands: North-Holland* (1990).
- [18] T. Bröcker and T. tom Dieck, “Representations of Compact Lie Groups,” *New York, USA: Springer Verlag* (1985).
- [19] A. Borel, *Natl. Acad. Sci.* **40** (1954) 1147.
- [20] M. Bando, T. Kuramoto, T. Maskawa and S. Uehara, *Prog. Theor. Phys.* **72** (1984) 313.
- [21] M. Bando, T. Kuramoto, T. Maskawa and S. Uehara, *Phys. Lett.* **138B** (1984) 94.
- [22] S. Coleman, J. Wess and B. Zumino, *Phys. Rev.* **177** (1969) 2239.
- [23] C.G. Callan, S. Coleman, J. Wess and B. Zumino, *Phys. Rev.* **177** (1969) 2247.
- [24] J.P. Serre *Séminaire Bourbaki, Exposé* **100** (1954) 1-8.
- [25] Y. Achiman, S. Aoyama, and J. W. van Holten, *Nucl. Phys.* **B258** (1985) 179.
- [26] C.-L. Ong, *Phys. Rev.* **D27** (1983) 911.
- [27] M. P. Mattis, *Phys. Rev.* **D28** (1983) 2649.
- [28] J. W. van Holten, *Z. Phys.* **C27** (1985) 57.